



FRUIT-TO-SIZE™

Is a concentrated foliar product which is specifically formulated to reduce stress and increase fruit size.

FOLIAR AND FRUIT ENHANCER

(N% - P% - K% - S%)
(4 - 8 - 24 - 5)

ACTIVE INGREDIENTS

Element	%W/W	g/kg	Element	%W/W	mg/kg
Nitrogen (N)	4.32	43	Manganese(Mn)	0.028	280
Phosphate (P)	8.37	84	Zinc (Zn)	0.255	2 550
Potassium (K)	23.75	232			
Magnesium (Mg)	3.51	35			
Sulphur (S)	5.04	50			

Batch number:

Date of manufacture:



Agri Technovation NZ Ltd
27A Wicklam Lane,
Greenhithe
Auckland 0632

info@agritechnovation.co.nz
www.agritechnovation.co.nz
Manufactured in New Zealand

Product information and crop recommendations provided on this label are only valid for New Zealand.



25KG ● 500KG ● 1000KG ●

FRUIT-TO-SIZE™

GENERAL INFORMATION

For optimum results FRUIT-TO-SIZE™ should be used in a plant nutrition programme. Suitable for use in combination with other nutrients, if directed by an Agri Technovation horticulturalist. Always contact your agronomist for crop specific information.

DIRECTIONS FOR USE

- Avoid application under extreme conditions, such as temperatures above 30°C, wind speeds exceeding 8 km/h or when plants are wilting.
- Wear suitable protective clothing when sprayed with other products.
- For best results with this product, the pH of the tank mixture should be between 5 and 6.
- This product should be applied as a foliar spray when crops have sufficient leaf area.
- Recommendations should be based on soil and leaf analyses.
- Where overhead irrigation systems are in use, avoid scheduling irrigation directly after application.
- In orchards equipped with micro irrigation systems, spray deposits should not be affected by irrigation.

MIXING

Always mix FRUIT-TO-SIZE™ with sufficient quantities of water. Never apply FRUIT-TO-SIZE™ undiluted. Follow standard tank mixing procedures. Ensure good agitation throughout application.

COMPATIBILITY

It is advisable to consult with a certified agrochemical specialist. Always perform a standard jar test prior to mixing to ensure compatibility of products to be used in tank mixture.

WARNING

Do not swallow. The dust from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin. Users are encouraged to seek further advice.

CAUTION

FRUIT-TO-SIZE™ is recommended for use on all crops during plant growth periods that require relatively high levels of nitrogen, phosphate, potassium, magnesium, sulfur, manganese and zinc. FRUIT-TO-SIZE™ is fully water soluble and provides readily available nutrients as shown in the guaranteed analysis for foliar applications.

FRUIT-TO-SIZE™ will be influenced by differences in crop varieties, soil and environmental conditions and should be applied under conditions of optimum plant growth. Optimum conditions are defined as adequate humidity, moderate temperatures and adequate soil moisture. Optimum conditions for foliar application to most crops generally occur during early morning or late evening hours.

CONDITIONS OF SALE

Every care is taken during the manufacturing of this product. However, as environmental conditions can vary considerably, no responsibility can be taken by the manufacturer, importer or his agents for any injury, damage, loss or any other detrimental effects following the use or misuse of this product.

Should this product not meet your expectations, please contact your fertiliser supplier before use. Before using this product seek appropriate agronomic advice.

Product information and crop recommendations provided on this label are only valid for New Zealand.

New Zealand Poison Hot-line number:
0800 764 766

APPLICATION RATES

Crop	Maximum Rate per Hectare*	Rate per 100 litre Water	Additional Information
Pome Fruit	15 kg	500 g	Apply 2 sprays the first 3 weeks in November, at 7-10 day intervals.
Citrus	20 kg	500 g	Apply 6 weeks after 100% petal fall, during the cell enlargement period. Repeat the same spray 4 weeks later and then according to fruit size and expected yield potential.
Wine Grapes	7 kg	500 g	Apply as a spray at 4 mm berry size and repeat at the start of veraison.
Nuts	20 kg	500 g	Apply as full cover spray when the nut diameter is between 18-23 mm and repeat the spray when the nut diameter is between 26-30 mm.
Avocados	20 kg	500 g	Apply 6 weeks after 100% petal fall, during the cell enlargement period. Repeat the same spray 4 weeks later and then according to fruit size and expected yield potential.
Stone Fruit	15 kg	500 g	Apply at pit hardening and repeat 14 days later.
Berries	4 kg	800 g	Apply during fruit fill and ripening at 10-14 day intervals as required.
Kiwifruit	7 kg	500g	Apply as a spray at 4mm berry size and repeat at the start of veraison.

* Maximum rates are dependent on crop size.

