

AVO-TO-GROWTM

Is a foliar fertiliser formulated specifically for avocados containing macro and chelated microelements.


FOLIAR SPRAY

(N% - P% - K% - S%)

(22 - 3 - 7 - 10)

NUTRIENT ANALYSIS

Element	%W/W	g/kg
Nitrogen (N)	2.86	219
Phosphate (P)	2.61	26
Potassium (K)	7.20	72
Magnesium (Mg)	0.77	8
Sulphur (S)	10.04	100





Agri Technovation NZ Ltd 27A Wicklam Lane, Greenhithe Auckland 0632

info@agritechnovation.co.nz www.agritechnovation.co.nz Manufactured in New Zealand

Product information and crop recommendations provided on this label are only valid for New Zealand.



AVO-TO-GROWTM

GENERAL INFORMATION

AVO-TO-GROW™ is a foliar formulated specifically for avocados containing macro and chelated micro elements. Consult with a certified horticulturist, agronomist or your crop advisor, consultant or the registration holder should crop specific programmes be required, or if any additional information or guidance is required concerning the use of this product. Suitable in combination with other nutrients. For optimum results AVO-TO-GROW™ should be used in combination with a fertiliser programme. Recommendation should be based on soil and leaf analyses.

DIRECTIONS FOR USE

- Do not apply to crops that are water or heat stressed or in very windy conditions.
- Wear suitable protective clothing when sprayed with other products.
- For best results with this product, the pH of the tank mixture should be between
 5 and 6
- This product should be applied as a foliar spray when crops have sufficient leaf area
- Where overhead systems are in use, avoid scheduling irrigation directly after application.
- In orchards equipped with micro irrigation systems, spray deposits should not be affected by irrigation.

MIXING

Always mix AVO-TO-GROW™ with sufficient quantities of water. Never apply AVO-TO-GROW™ undiluted. Follow standard tank mixing procedures. Ensure good agitation throughout application.

COMPATIBILITY

It is advisable to consult with a certified agrochemical specialist. Always perform a standard jar test prior to mixing to ensure compatibility of products to be used in tank mixture.

WARNING

Do not swallow. The dust from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin. Users are encouraged to seek further advice.

CAUTION

Do not store in direct sunlight. Store the product inside at room temperature. This is a blended fertiliser in which some segregation of the ingredients may occur that may lead to some variation from the stated analysis.

CONDITIONS OF SALE

Every care is taken during the manufacturing of this product. However, as environmental conditions can vary considerably, no responsibility can be taken by the manufacturer, importer or his agents for any injury, damage, loss or any other detrimental effects following the use or misuse of this product.

Should this product not meet your expectations, please contact your fertiliser supplier before use. Before using this product seek appropriate agronomic advice.

Product information and crop recommendations provided on this label are only valid for New Zealand.

New Zealand Poison Hot-line number 0800 764 766

APPLICATION RATES

Growth Stage	g/100 liters of water	kg/ha (Max application rate)*	
Bearing trees	500 g	15 kg	
First Spray: Post harvest depending on cultivar.			
Second Spray: At budburst.			
Third Spray: After second fruit drop.			

^{*} Maximum rates are dependent on crop size.