

AMINOPRO™

NATURAL BIOSTIMULANT TO SUPPORT PLANT METABOLISM

(N% - P% - K% - S%)
(1 - 0 - 0 - 0)

NUTRIENT ANALYSIS

Element	%W/V	g/l
Nitrogen (N)	1.2	12.0

pH = 3.8 ± 0.2

SG (@ 20°C) = 1.04 ± 0.02



Agri Technovation Ltd
27A Wicklam Lane,
Greenhithe
Auckland 0632
09 954 5411
info@agritechnovation.co.nz
www.agritechnovation.co.nz
Manufactured in New Zealand
New Zealand Poison Hot-line number: 0800 764 766



Batch number:

Date of manufacture:



AMINOPRO™

GENERAL INFORMATION

AMINOPRO™ is an organically based liquid biostimulant to promote plant growth and support recovery. Use AMINOPRO™ as advised by your qualified agricultural advisor. Directly contact your qualified agricultural advisor for any additional information or guidance required concerning the use of this product. For optimum results AMINOPRO™ should be used in combination with a fertiliser programme.

DIRECTIONS FOR USE

- For foliar application
- Avoid application under extreme conditions such as temperatures above 30 °C, wind speeds exceeding 15 km/h and wilting conditions
- For improved uptake, spray in the early morning or evening, avoid spraying in the heat of the day
- Suitable for use in combination with other nutrients
- Wear suitable protective clothing
- Keep container closed when not in use

COMPATIBILITY

It is advisable to consult with a certified agrochemical specialist. It is required to perform a miscibility test prior to mixing with other chemicals.

However do NOT apply AMINOPRO™:

- In a mix with Amine herbicides
- In combination with copper containing fungicides, sulphates, phosphates or phosphonates
- Within 14 days before or after a copper application
- In combination with highly alkaline water sources and/or materials

User assumes full responsibility to ensure compatibility when tank mixing with other products. Water quality testing is advisable to ensure full compatibility and optimal product performance.

MIXING

Shake well before use. Always mix AMINOPRO™ with sufficient quantities of water. Never apply AMINOPRO™ undiluted. Follow standard tank mixing procedures. Ensure good agitation throughout application. Do not store mixed product.

WARNING

Avoid contact with eyes. Keep out of reach of children. Do not swallow. If swallowed seek medical advice. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin. Users are encouraged to seek further advice. Water quality can affect this product's performance. Water quality and pH testing needs to be conducted to ensure optimal product performance. Water quality improvement products may be required to adjust water quality parameters before mixing with AMINOPRO™.

CAUTION

Do not store in direct sunlight. Store the product inside at room temperature. This is a blended fertiliser in which some segregation of the ingredients may occur that may lead to some variation from the stated analysis.

CONDITIONS OF SALE

Although this product has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic conditions as well as method, time and accuracy of application. Every care is taken during the manufacture of this product. However, as environmental conditions can vary considerably, the manufacturer, importer and/or its agents do not accept any responsibility for any injury, damage, loss, and/or any other detrimental effects or consequences following the misuse of this product.

APPLICATION RATES

Crop	Maximum Rate per Hectare*	Rate per 100 ℓ Water
Grains	1-2 ℓ/ha	250 - 500 mℓ
Vegetables	2 - 3 ℓ/ha	500 - 1 000 mℓ
Fruit trees	3 - 4 ℓ/ha	150 - 200 mℓ

*Maximum rates are dependent on crop size