

KIWI STRENGTH™

FOLIAR FERTILISER FORMULATED TO ENHANCE CELL WALL STRENGTH

(N% - P% - K% - S%)
(1 - 0 - 0 - 0)

NUTRIENT ANALYSIS

Element	%W/V	g/l
Nitrogen (N)	0.50	5.00
Calcium (Ca)	2.30	23.00
Magnesium (Mg)	0.40	4.00

pH = 5.7 ± 0.2

SG (@ 20°C) = 1.08 ± 0.02

Element	%W/V	mg/l
Boron (B)	1.080	10.80
Silicon (Si)	0.028	0.28



Batch number:

Date of manufacture:



Agri Technovation Ltd
27A Wicklam Lane,
Greenhithe
Auckland 0632
09 954 5411

info@agritechnovation.co.nz
www.agritechnovation.co.nz
Manufactured in New Zealand
New Zealand Poison Hot-line number: 0800 764 766



KIWI STRENGTH™

GENERAL INFORMATION

KIWI STRENGTH™ is a calcium, boron and orthosilicic acid fertiliser, blended to support plant growth. Use KIWI STRENGTH™ as advised by your qualified agricultural advisor. Directly contact your qualified agricultural advisor for any additional information or guidance required concerning the use of this product. For optimum results KIWI STRENGTH™ should be used in combination with a fertiliser programme.

DIRECTIONS FOR USE

- To get optimum results with this product, make sure the pH value of the mixture in the tank is between 5-6
- This product should be applied as a foliar spray when crops have sufficient leaf area
- Do not exceed the appropriate application rates
- Avoid application under extreme conditions such as temperatures above 30 °C, wind speed exceeding 15 km/h or wilting conditions
- For superior uptake, spray in the early morning or evening, avoid spraying in the heat of the day
- Always contact your agriculturalist for crop specific information
- Wear suitable protective clothing.
- **FOLIAR APPLICATIONS:** Do not exceed a 1% solution (One part KIWI STRENGTH™ to 100 parts water)
- Keep container closed when not in use

COMPATIBILITY

It is advisable to consult with a certified agrochemical specialist. It is required to perform a miscibility test prior to mixing with other chemicals.

However do NOT apply KIWI STRENGTH™:

- In a mix with Amine herbicides
- In combination with copper containing fungicides, sulphates, phosphates or phosphonates
- Within 14 days before or after a copper application
- In combination with highly alkaline water sources and/ or materials
- With water sources or products containing, carbonates, sulphur or phosphate

User assumes full responsibility to ensure compatibility when tank mixing with other products. Water quality testing is advisable to ensure full compatibility and optimal product performance.

MIXING

Always mix KIWI STRENGTH™ with sufficient quantities of water. Follow standard tank mixing procedures. Ensure good agitation throughout application. Do not store mixed product.

WARNING

Avoid contact with eyes. Keep out of reach of children. Do not swallow. If swallowed seek medical advice. The dust from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin. Users are encouraged to seek further advice.

CAUTION

Do not store in direct sunlight. Store the product inside at room temperature. This is a blended fertiliser in which some segregation of the ingredients may occur that may lead to some variation from the stated analysis.

CONDITIONS OF SALE

Although this product has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic conditions as well as method, time and accuracy of application. Every care is taken during the manufacture of this product. However, as environmental conditions can vary considerably, the manufacturer, importer and/or its agents do not accept any responsibility for any injury, damage, loss, and/or any other detrimental effects or consequences following the misuse of this product.

APPLICATION RATES

Crop	Maximum Rate per Hectare*	Rate per 100 litre Water	Additional Information
Kiwifruit	5 l	500ml/100 l	Do not mix with copper, apply twice pre-flower and twice post-harvest

*Maximum rates are dependent on crop size.