

# CITRUS-TO-GROW™

## FOLIAR FERTILISER

( N% - P% - K% - S% )

( 5 - 17 - 8 - 5 )



### NUTRIENT ANALYSIS

Element	%W/W	g/kg	Element	%W/W	mg/kg
Nitrogen (N)	4.8	48	Zinc (Zn)	0.229	2 290
Phosphate (P)	17.0	170	Boron (B)	0.147	1 470
Potassium (K)	7.6	76	Manganese (Mn)	0.143	1 350
Sulphur (S)	4.9	49	Iron (Fe)	0.100	1 000
Magnesium (Mg)	3.4	34	Molybdenum	0.027	270

### GENERAL INFORMATION

CITRUS-TO-GROW™ highly soluble, multi elemental fertiliser. Use CITRUS-TO-GROW™ as advised by your qualified agricultural advisor. Directly contact your qualified agricultural advisor for any additional information or guidance required concerning the use of this product. For optimum results CITRUS-TO-GROW™ should be used in combination with a fertiliser programme.

### DIRECTIONS FOR USE

- To get optimum results with this product, make sure the pH value of the mixture in the tank is between 5-6
- Do not exceed the appropriate application rates
- Avoid application under extreme conditions such as temperatures above 30 °C, wind speed exceeding 15 km/h or wilting conditions
- Wear suitable protective clothing
- Clean equipment before and after use
- Do not apply to fruit after color break, only apply to fruit before color break (green fruit)

### COMPATIBILITY

It is advisable to consult with a certified agrochemical specialist. It is required to perform a miscibility test prior to mixing with other chemicals.

However do NOT apply CITRUS-TO-GROW™:

- In a mix with Amine herbicides
- In combination with copper containing fungicides, sulphates, phosphates or phosphonates
- Within 14 days before or after a copper application
- In combination with highly alkaline water sources and/ or materials
- With water sources or products containing high levels of calcium, magnesium and/or carbonates

User assumes full responsibility to ensure compatibility when tank mixing with other products. Water quality testing is advisable to ensure full compatibility and optimal product performance.

### MIXING

Always mix CITRUS-TO-GROW™ with sufficient quantities of water. Follow standard tank mixing procedures. Ensure good agitation throughout application. Do not store mixed product.

### WARNING

Avoid contact with eyes. Keep out of reach of children. Do not swallow. If swallowed seek medical advice. The dust from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin. Users are encouraged to seek further advice.

### CAUTION

Do not store in direct sunlight. Store the product inside at room temperature. This is a blended fertiliser in which some segregation of the ingredients may occur that may lead to some variation from the stated analysis.

### CONDITIONS OF SALE

Although this product has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic conditions as well as method, time and accuracy of application. Every care is taken during the manufacture of this product. However, as environmental conditions can vary considerably, the manufacturer, importer and/ or its agents do not accept any responsibility for any injury, damage, loss, and/or any other detrimental effects or consequences following the misuse of this product.

### APPLICATION RATES

Growth Stage	Rate per 100 ℓ water	Maximum Rate per Hectare
Bearing trees	500g	15kg
First Spray: Pre-and post harvest (around March, April and May)		
Second Spray: At budburst		
Third Spray: End January		

\* Maximum rates are dependent on crop size.



Agri Technovation Ltd  
27A Wicklam Lane,  
Greenhithe  
Auckland 0632  
09 954 5411  
[info@agrитеchnovation.co.nz](mailto:info@agrитеchnovation.co.nz)  
[www.agritechnovation.co.nz](http://www.agritechnovation.co.nz)  
Manufactured in New Zealand  
New Zealand Poison Hot-line number: 0800 764 766



Batch number:

Date of manufacture:

