

FRUIT-TO-SIZE™

WATER-SOLUBLE FOLIAR FERTILISER

(N% - P% - K% - S%)

(4 - 8 - 23 - 5)

NUTRIENT ANALYSIS

Element	%W/W	g/kg	Element	%W/W	mg/kg
Nitrogen (N)	4.3	43	Manganese(Mn)	0.028	280
Phosphate (P)	8.4	84	Zinc (Zn)	0.255	2 550
Potassium (K)	23.2	232			
Magnesium (Mg)	3.5	35			
Sulphur (S)	5.0	50			

GENERAL INFORMATION

FRUIT-TO-SIZE™ is a water-soluble foliar fertiliser containing macro- and micronutrients to support fruit sizing. Use FRUIT-TO-SIZE™ as advised by your qualified agricultural advisor. Directly contact your qualified agricultural advisor for any additional information or guidance required concerning the use of this product. For optimum results FRUIT-TO-SIZE™ should be used in combination with a fertiliser programme.

DIRECTIONS FOR USE

- Avoid applications under extreme conditions, such as temperatures above 30°C, wind speeds exceeding 15 km/h or when plants are wilting
- For improved uptake, apply early in the morning or evening, and avoid application during the heat of the day
- Wear suitable protective clothing
- Optimum results with this product are obtained if the pH of the tank mix is between 5 and 6

COMPATIBILITY

It is advisable to consult with a certified agrochemical specialist. It is required to perform a miscibility test prior to mixing with other chemicals.

However do NOT apply FRUIT-TO-SIZE™:

- In a mix with Amine herbicides
- In combination with copper containing fungicides, sulphates, phosphates or phosphonates
- Within 14 days before or after a copper application
- In combination with highly alkaline water sources and/ or materials
- With water sources or products containing calcium and/or carbonates

User assumes full responsibility to ensure compatibility when tank mixing with other products. Water quality testing is advisable to ensure full compatibility and optimal product performance.

MIXING

Always mix FRUIT-TO-SIZE™ with sufficient quantities of water. Follow standard tank mixing procedures. Ensure good agitation throughout application. Do not store mixed product.

WARNING

Avoid contact with eyes. Keep out of reach of children. Do not swallow. If swallowed seek medical advice. The dust from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin.

CAUTION

Do not store in direct sunlight. Store the product inside at room temperature. This is a blended fertiliser in which some segregation of the ingredients may occur that may lead to some variation from the stated analysis.

CONDITIONS OF SALE

Although this product has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic conditions as well as method, time and accuracy of application. Every care is taken during the manufacture of this product. However, as environmental conditions can vary considerably, the manufacturer, importer and/or its agents do not accept any responsibility for any injury, damage, loss, and/or any other detrimental effects or consequences following the misuse of this product.

APPLICATION RATES

Crop	Maximum Rate per Hectare*	Rate per 100 litre Water	Additional Information
Pome fruit	15 kg	500 g	Apply as a spray 40 days after full bloom.
Citrus	20 kg	500 g	Apply 6 weeks after 100% petal fall, during the cell enlargement period. Repeat the same spray 4 weeks later and then according to fruit size and expected yield potential.
Wine Grapes	7 kg	500 g	Apply as a spray at 4 mm berry size and repeat at the start of veraison.
Nuts	20 kg	500 g	Apply as full cover spray when the nut diameter is between 18-23 mm and repeat the spray when the nut diameter is between 26-30 mm.
Avocados	20 kg	500 g	Apply 6 weeks after 100% petal fall, during the cell enlargement period. Repeat the same spray 4 weeks later and then according to fruit size and expected yield potential.
Stone Fruit	15 kg	500 g	Apply at pit hardening and repeat 14 days later.
Berries	4 kg	800 g	Apply during fruit fill and ripening at 10-14 day intervals as required.
Kiwifruit	7 kg	500 g	Apply as a spray at 4mm berry size and repeat at the start of veraison.

*Maximum rates are dependent on crop size.



Agri Technovation Ltd
27A Wicklam Lane,
Greenhithe
Auckland 0632
09 954 5411

info@agritechnovation.co.nz
www.agritechnovation.co.nz
Manufactured in New Zealand

New Zealand Poison Hot-line number: 0800 764 766



Batch number:

Date of manufacture:



25 KG

500 KG

1000 KG