

GENERAL INFORMATION

POLLINATOR™ is a product formulated to support pollination. Use POLLINATOR™ as advised by your qualified agricultural advisor. Directly contact your qualified agricultural advisor for any additional information or guidance required concerning the use of this product. For optimum results POLLINATOR™ should be used in combination with optimum agricultural crop improvement practices.

DIRECTIONS FOR USE

- Avoid applications under extreme conditions, such as temperatures above 30°C, wind speeds exceeding 15 km/h or when plants are wilting
- For improved uptake, apply early in the morning or evening, and avoid application during the heat of the day
- Wear suitable protective clothing
- Optimum results with this product are obtained if the pH of the tank mix is between 5 and 6
- Keep container closed when not in use
- Spray after pollen is distributed

COMPATIBILITY

It is advisable to consult with a qualified agricultural advisor. It is required to do a miscibility test prior to mixing with other chemicals.

However do NOT apply POLLINATOR™:

- In a mix with amine herbicides
- In combination with copper containing fungicides, sulphates, phosphates or phosphonates
- Do not apply Pollinator 3 days before a copper fungicide application or 10 days after a copper fungicide application

User assumes full responsibility to ensure compatibility when tank mixing with other products. Water quality testing is advisable to ensure full compatibility and optimal product performance.

MIXING

Shake well before use. Never apply POLLINATOR™ undiluted. Follow standard tank mixing procedures. Ensure good agitation throughout application. Do not store mixed product.

WARNING

Avoid contact with eyes. Keep out of reach of children. Do not swallow. If swallowed seek medical advice. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin. Users are encouraged to seek further advice.

CAUTION

Do not store in direct sunlight. Store the product inside at room temperature. This is a blended fertiliser in which some segregation of the ingredients may occur that may lead to some variation from the stated analysis.

Pollinator™

PRODUCT TO SUPPORT POLLINATION

(N% - P% - K% - S%)
(0 - 1 - 0 - 0)

ACTIVE INGREDIENTS

Element	% W/V	g/ℓ
Phosphorus (P)	0.52	5.20
Calcium (Ca)	0.52	5.20

pH = 3.7 ± 0.2

SG (@ 20°C) = 1.05 ± 0.02

Batch number:

Date of manufacture:

Agri Technovation Ltd
27A Wicklam Lane,
Greenhithe Auckland 0632
09 954 5411
info@agritechnovation.co.nz
www.agritechnovation.co.nz
Manufactured in Australia
New Zealand Poison Hot-line number: 0800 764 766



CONDITIONS OF SALE

Although this product has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic conditions as well as method, time and accuracy of application. Every care is taken during the manufacture of this product. However, as environmental conditions can vary considerably, the manufacturer, importer and/or its agents do not accept any responsibility for any injury, damage, loss, and/or any other detrimental effects or consequences following the misuse of this product.

APPLICATION RATES

Crop	Maximum Rate per Hectare*	Additional Information
Kiwi Fruit	3 ℓ	Apply at flowering, after peak pollination window to coincide with bee activity
Citrus	3 - 4 ℓ	Apply product at the onset of flowering (approximately 30% flowering) and repeat product application at full bloom
Pome- & Stone Fruit	3 - 4 ℓ	Apply at first/ full bloom
Blueberries	3 - 4 ℓ	Apply every 2 weeks during flowering
Tree Nuts	3 - 4 ℓ	Apply at open flower. Maximum three applications
Soybeans	3 - 4 ℓ	Apply at R1/R2
Oilseed Crops	3 - 4 ℓ	Sunflower: Apply at early/full flowering Canola: Apply at early/ full flowering
Table and Wine grapes	3 - 4 ℓ	Apply when 20% of flowerhoods have fallen, and again at full bloom
Wheat/Barley	3 - 4 ℓ	Apply at flowering
Maize	3 - 4 ℓ	Apply product at onset of flowering and do a repeat application at full bloom (VT-R1)
Seed Maize	2 ℓ	2 applications, one at 50% silking and one at 80% silking
Avocados	3 - 4 ℓ	Apply at 25% flower, repeat at 75% - 100% flower. Apply after pollination on female flowers.
Tomato	3 - 4 ℓ	Apply product at onset of flowering and again full bloom
Dry Beans	2 ℓ	For dry beans: 2 applications, one at R1 (first flower) and one at R3 (full flower)

*Maximum rates are dependent on tree size.



1ℓ