

# Kelpura™

HIGHLY CONCENTRATED PURE LIQUID KELP

pH =  $3.8 \pm 0.2$  | SG =  $1.02 \pm 0.02$


( N% - P% - K% - S% )

( 0 - 0 - 0 - 0 )



Controlled by ECOCERT SAS

Product suitable for use in Organic Farming in accordance with EU Regulations 2018/848 and 2021/1165



20L

200L

1000L



Technovation

Specialized to succeed

Agri Technovation Ltd  
27A Wicklam Lane,  
Greenhithe  
Auckland 0632

09 954 5411

info@agritechnovation.co.nz

www.agritechnovation.co.nz

Manufactured in South Africa

New Zealand Poison


Hot-line number: 0800 764 766



Batch number:



Date of manufacture:





## GENERAL INFORMATION

KELPURA™ is a pure kelp extract derived from *Ecklonia maxima*. Use KELPURA™ as advised by your qualified agricultural advisor. Directly contact your qualified agricultural advisor for any additional information or guidance required concerning the use of this product. For optimum results KELPURA™ should be used in combination with a optimum agricultural crop improvement practices.

## DIRECTIONS FOR USE

- Avoid applications under extreme conditions, such as temperatures above 30°C, wind speeds exceeding 15 km/h or when plants are wilting
- For improved uptake, spray in the early morning or evening, avoid spraying during the heat of the day.
- Wear suitable protective clothing when sprayed with other products
- Do not apply through filters smaller than 100 micron
- **Foliar:** Suitable for use with all forms of foliar application systems. For best results, apply when plants have sufficient leaf area. Dilute 1 part KELPURA™ concentrate with 200 parts of water. Apply 2 - 5 ℓ per hectare. To get optimum results with this product make sure the pH of the mixture in the tank is between 5 - 6.

## COMPATIBILITY

It is advisable to consult with a qualified agricultural advisor. It is required to perform a miscibility test prior to mixing with other chemicals.

However do NOT apply KELPURA™:

- In a mix with amine herbicides
- In combination with copper containing fungicides, sulphates, phosphates or phosphonates
- Within 14 days before or after a copper application
- In combination with highly alkaline water sources and/or materials

User assumes full responsibility to ensure compatibility when tank mixing with other products. Water quality testing is advisable to ensure full compatibility and optimal product performance.

## MIXING

Shake well before use. Always mix KELPURA™ with sufficient quantities of water. Never apply KELPURA™ undiluted. Follow standard tank mixing procedures. Do not store mixed product

## WARNING

Avoid contact with eyes. Keep out of reach of children. Do not swallow. If swallowed seek medical advice. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin.

## CAUTION

Avoid contact with eyes. Keep out of reach of children. If swallowed seek medical advice. Do not store in direct sunlight. Store product inside at room temperature

## CONDITIONS OF SALE

Although this product has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic conditions as well as method, time and accuracy of application. Every care is taken during the manufacture of this product. However, as environmental conditions can vary considerably, the manufacturer, importer and/or its agents do not accept any responsibility for any injury, damage, loss, and/ or any other detrimental effects or consequences following the misuse of this product.

## APPLICATION RATES

| Application        | *Maximum Rate per Hectare | Additional Information                                                                       |
|--------------------|---------------------------|----------------------------------------------------------------------------------------------|
| Foliar application | 2 - 5 ℓ                   | Optimum results with this product are obtained if the pH of the tank mix is between 5 and 6. |

\* Maximum rates are dependent on crop size.

